



## **COVID-19 Guidance for Licensed and Group Child Care Homes**

February 18, 2022

### Introduction

Johnson County Department of Health and Environment's childcare recommendations reflect the current understanding of COVID-19 transmission in children. As new evidence emerges, guidance may change.

On December 27th, 2021, the Centers for Disease Control and Prevention released updated recommendations regarding COVID-19 isolation and quarantine measures for the general public. While recognizing people may remain infectious (capable of spreading COVID-19) for up to 10 days, current data indicates that most people are infectious early in their infection. As a result, CDC shortened the recommended isolation period for infected individuals to five days if they are asymptomatic or their symptoms are resolving (without fever for 24 hours), followed by five days of mask wearing when around others.

Due to the unique constraints of the childcare home (e.g., limited space to separate positive children, etc.), JCDHE recognizes that there may be some additional risk of transmission with the implementation of the new guidance. As such, JCDHE will support childcare providers in implementing the shortened isolation protocol for those over the age of two that are willing and able to mask as well as those providers who choose to stay with the previous guidance.

### Definitions

**Quarantine:** Keeps someone who might have been exposed to the virus away from others. Individuals in quarantine should stay home. If an individual must be in public to seek medical assistance, they should practice masking and physical distancing as much as possible. Quarantine/exclusion timelines always begin with the last exposure to a person with confirmed or presumed COVID-19. Guidance from CDC and the Kansas Department of Health and Environment also offers the following shortened quarantine protocols.

#### **For individuals who are willing and able to mask:**

- Five-day quarantine followed by five days of consistent mask wearing while around others, as long as the individual remains symptom-free. Testing on day five after exposure is

strongly recommended.

**For individuals who are unwilling or unable to mask (e.g., under the age of two):**

- Option 1 (with testing): Exposed individuals who remain asymptomatic may receive a test on or after day six. If there is a negative result and no symptoms, individuals may return to normal activities on day eight after exposure.
- Option 2 (without testing): Individuals who are exposed and remain asymptomatic, but do NOT take a test, should quarantine for 10 days, returning to activities on day 11 after exposure. Although the risk of transmission after the 10-day quarantine period is low, the risk is not zero.

Childcare facilities may stay with the 10-day quarantine as it offers the most protection to the facility. The facility is able to cohort if there is only one case identified (discussed in later section).

**Isolation:** Isolation separates people who are infected with the virus away from people who are not infected. Individuals with confirmed or presumed COVID-19 should isolate within their household and use a separate bedroom/bathroom. Sleeping areas should not be shared. Individuals should not spend time in common household areas (living room, kitchen); if face-to-face interactions must take place, all household members should mask. Disinfect frequently touched surfaces in the household often.

**Close Contact/Exposure:** A close contact is defined as:

- a. Being directly exposed to infectious secretions (e.g., being coughed on); or
- b. Being within 6 feet for 15 cumulative minutes or more over a 24-hour period. Additional factors like infected person/contact masking (i.e., both the infectious individual and the potential close contact have been consistently and properly masked), classroom-level mitigation measures, individual risk profiles, and case symptomology may affect this determination.

Either (a) or (b) is defined as close contact if it occurred during the case's infectious period.

**Infectious Period:** An individual is considered **infectious** (capable of spreading the virus) for two days *before* their symptoms began until ten days *after* symptom onset *and* 24 hours after their fever (if present) has resolved without the aid of medication *and* initial symptoms have improved. For an **asymptomatic** individual who tests positive for COVID-19, their infectious period is two days before through 10 days after their specimen was collected.

**Presumed Positive:** Symptomatic individuals with a known exposure to a COVID-19 positive individual within the 14 days prior to symptom onset are presumed positive. Becoming symptomatic while excluded for quarantine should trigger a move from quarantine to isolation.

**Vaccine (COVID-19) Breakthrough Case:** A breakthrough case is defined as an individual who has a laboratory detected on a respiratory specimen collected greater than or equal to 14 days after completing the FDA-authorized COVID-19 vaccine.

**Screening:** Screening remains in place because the virus is still circulating. Screen children and staff daily before admittance for signs and symptoms of illness. Screening includes asking questions, observing for signs of illness and *could* include checking for fever if child appears ill. Many facilities are doing after nap temperatures.

- Has there been an exposure to someone diagnosed with COVID-19, either household or non-household contact?
- Is anyone in the home showing signs of illness or who have the following:
  - o fever greater than 100.4 degrees (F) (need to be aware of person's "normal" temperature as some people run lower "normal" and therefore a fever for them could lower than 100.4)
  - o cough
  - o shortness of breath/difficulty breathing
  - o sudden loss of smell and/or taste (not as common with Omicron)
  - o other signs of illness (headache, sore throat, general aches/pains, fatigue/weakness/extreme exhaustion)
  - o Check the child's temperature as indicated. Per childcare regulation, sick children should not be in the childcare setting.

**Masks:** Please see CDC guidance for appropriate types of masks:

<https://www.cdc.gov/coronavirus/2019ncov/preventgetting-sick/about-face-coverings.html>

The CDC recommends the most protective mask that fits well (i.e., fitting closely on the face without any gaps along the edges or around the nose) and can be comfortably worn throughout the day.

**Recommended Masks:**

- Respirator (N95)
- KN95
- Medical or surgical masks
- Properly fitting masks (i.e., snugly around nose AND chin with no gaps around the sides of the face)
- Masks made with tightly woven fabric (i.e., fabrics that do NOT let light pass through when held up to a light source)
- Masks with two or three layers
- Masks with inner filter pockets

## Guidance for Operations

### Parents drop off and pick up

- JCDHE STRONGLY recommends that all parents, visitors and others entering the daycare home are masked no matter their vaccination status.
- Provider should still limit access to the day care home as much as possible and may want to consider door/porch drop off versus allowing parents inside.

### Bringing in outside resources/activities

Many home providers bring in outside activities instead of taking all the children to various locations. While it is best to continue to limit who comes into the home, if the provider chooses to bring in activities, the person should be vaccinated, wear an appropriate mask over mouth and nose at all times (preferably an N95 or KN95) and maintain as much physical distancing as possible.

### Sensory tables and supplies

CDC guidance has not been updated regarding this issue. However, if children wash their hands prior to and after usage, JCDHE feels it is safe to restart sensory table usage. Capacity at a sensory table should be limited to one or two children at a time depending on size. Individual supplies such as markers, crayons, and playdoh – should follow the same guidelines as sensory tables regarding hand washing and number of children in one location. Remember to clean and disinfect all toys, supplies and surfaces on a routine basis.

### Field trips

For those too young to be in school field trips are not recommended at this time due to the large amount of community spread.

### Mask usage

#### **Providers**

JCDHE **strongly recommends** that providers wear masks during childcare hours regardless of vaccine status, however you do not have to be masked when outside. For unvaccinated providers, it is recommended they receive the vaccine as soon as possible (if eligible) and should always wear a mask unless outside or physically distanced. For vaccinated providers, the data show the vaccine is working against severe illness; however, infections in vaccinated individuals can occur. For vaccinated individuals, masks are an extra layer of protection against infection, especially while community transmission is elevated.

Snug fitting masks with a see-through face area are acceptable. JCDHE does not approve face shields alone. Masks should be worn at least at children's pick up and drop off times and when other adult non-residents are in the home. This is especially important if parents are coming into the childcare home.

Since children are unmasked during mealtime, it is recommended that providers not eat lunch with the children due to the added risk of exposure. When providers are at least six feet from others, such as bathroom or at nap time, their mask may be removed for a break. Supervision must be maintained at all times.

When childcare surveyors enter a childcare home, there will be a written consultation on the NOSF regarding whether masks are worn by the provider and children.

### **Children**

In alignment with CDC and American Academy of Pediatrics, JCDHE strongly recommends that children over the age of two should be encouraged to wear masks when inside. The masks need to be removed for nap time and meals. Physical distancing of at least six feet is strongly encouraged for meal and nap time. Children should be spaced apart as much as possible during mealtime. The childcare home must also plan for what to do with the masks at mealtime to keep them from becoming soiled or contaminated, as well as a plan for changing out wet or otherwise contaminated children’s masks.

## **Management of Symptomatic Individuals in a Childcare Setting**

JCDHE recommends all licensed childcare facilities follow the below guidance for exclusion criteria and management of symptomatic individuals.

Individuals who meet the below criteria should be encouraged to seek testing for COVID-19. If a physician indicates the symptoms are due to a different *non-infectious* diagnosis (e.g., allergies, asthma), a child can be readmitted to childcare prior to improvement of their symptoms.

### **Primary symptoms (at least one)**

- Cough
- Shortness of breath
- Difficulty breathing
- Loss of taste and/or smell

### **Secondary Symptoms (at least two)**

- Fever (measured or subjective)
- Chills
- Muscle or body aches
- Headache
- Sore throat
- Diarrhea/nausea/vomiting
- Congestion/runny nose

Individuals who test negative for COVID-19 may return to childcare 24 hours after their symptoms improve.

## **Management of a COVID-19 Positive Individual**

### **Exclusion of new positive cases**

All individuals who test positive **must** be excluded from childcare settings for at least five days from when their symptoms began (or five days from the positive test if the individual doesn’t have

symptoms). An individual is considered **infectious** (capable of spreading the virus) two days before their symptoms began until 10 days after their symptom onset *and* 24 hours after their fever (if present) has resolved without the aid of medication *and* their initial symptoms have improved.

**For individuals who are willing and able to mask:**

**With symptoms:**

Isolate until:

- At least five days have passed since symptoms first appeared; AND
- At least 24 hours fever-free without the use of fever-reducing medications; AND
- Improvement in initial symptoms.

Mask for:

- Five additional days (day 6 through 10) using a high-quality mask when around others.

Testing recommendations:

- The **safest option** is to perform an antigen test on or around day five.
  - If negative, individual may return to childcare, but should continue to mask.
  - **If positive, individual should remain isolated until they have a negative test or have completed their 10-day isolation.**

**Without symptoms:**

Isolate for:

- Five days from day of test.

Mask for:

- Five additional days (day 6 through 10) using a high-quality mask when around others.

Testing recommendations:

- The **safest option** is to perform an antigen test on or around day five.
  - If negative, individual may return to childcare, but should continue to mask.
  - If positive, individual should remain isolated until they have a negative test or through day 10.

**For individuals who are unwilling or unable to mask (e.g., under the age of two), the safest option is a 10-day isolation period:**

**With symptoms:**

Isolate for:

- At least 10 days have passed since symptoms first appeared; AND
- At least 24 hours fever-free without the use of fever-reducing medications; AND
- Improvement in initial symptoms.

**Without symptoms:**

- Isolate for 10 days from a positive test. Use the date specimen was collected, not the

date of results.

If an individual is excluded with symptoms but the test comes back positive a few days later, the ten days is still based off the start of symptoms. If the individual is asymptomatic (not showing any symptoms), then the infectious period is two days before the date their lab test was collected until ten days after their lab test.

Please call JCDHE's Child Care Licensing hotline 913-477- 8361 (M-F, 8 a.m. – 5 p.m.) to report positive cases or exposures occurring in childcare –spreadsheet completion and submission is still needed.

### **Exclusion of Contacts**

#### **Contacts of a COVID-19 family member**

If a household member (e.g., spouse, kids, etc.) tests positive for COVID-19, then it is recommended that all unvaccinated household members quarantine following their last interaction with the positive case. If the positive individual can isolate in a separate bedroom, with a separate bathroom, spending little to no time in common areas, **NO** time in childcare areas and always wearing a mask in the presence of other household members, then the quarantine begins on the day the positive individual began isolating away from the household. Please contact JCHDE's Childcare Licensing Division to determine if the facility can remain open.

If it is not possible for the positive family member to isolate, household members will need to quarantine per current public health recommendations following the end of the infected person's isolation, and it may be recommended to close the childcare home. If additional household members become symptomatic/test positive during the isolation or quarantine period, the quarantine period starts over.

All close contacts, regardless of which quarantine procedure they follow, should continue to self-monitor for 10 days from exposure. If symptoms develop during the 10-day period, the person should self-isolate and get a PCR test.

#### **Contacts of a COVID-19 positive individual in a childcare home**

The choices are to:

- Cohort contacts of the positive child
- Close the childcare home

#### **Cohorting of the classroom**

The provider must notify all families of having a COVID-19 case. The provider can decide to remain open - however provider and residents are strongly encouraged to wear a mask except for eating. School age children are **strongly recommended** to wear masks. No field trips or other exposure to other people should occur. If the facility is on the food program – please notify the program so they can take

necessary precautions or move the visit. The childcare provider and children should not interact with any other children or have any outside events brought in (music, art, etc.) until 10 days from the last exposure to an infectious case.

#### Closing the childcare home

If only one positive child case is in the home, providers can decide to close the childcare home if they are more comfortable with this option. **If there are two or more positives in a childcare home**, JCDHE should be consulted to determine if there is evidence of COVID-19 transmission (two positives within 10 days of one another, without another known exposure). If internal transmission is identified, the recommendation may be that the entire facility be closed for up to 10 days from the last exposure to the infectious case.

If the childcare home chooses to close in lieu of cohorting, the return/reopen date is determined by one of the below options:

- Exposed persons who are able and willing to mask may return after day five provided they mask consistently around others for day six through day ten.  
OR
- Exposed persons testing negative on day six or later may return on day eight.  
OR
- the exposed persons who can't or chooses not to test and they remain asymptomatic through day 10, they can return on day 11.

#### **Notification following a COVID-19 positive (updated January 2022)**

When there is a positive in the classroom, families and staff must be notified of the positive case, while protecting confidentiality of the positive person's identity and encouraging them to monitor for signs and symptoms of COVID-19 for 10 days. It is important to communicate to families that even though they are allowed to cohort at the childcare facility, when not in the childcare they should remain at home for the duration of the recommended quarantine period. They should also be provided the guidance that was given by JCDHE to the provider. Template letters for providers have been sent out via email and are available at <https://www.jocogov.org/department/health/covid-19/child-care-guidance>.

#### **Post-exposure management of individuals with a lab confirmed positive case of COVID 19 in last 90 days (updated January 2022)**

Close contacts with evidence of previous infection **within the past 90 days** that is documented by a positive PCR or antigen test may be exempt from quarantine exclusion if they remain **asymptomatic** following their exposure. Positive serology or antibody tests may not be substituted for either the PCR or antigen test. If the close contact becomes symptomatic following their exposure, but it is during the 90 days after recovery from a prior infection, there is a possibility of reinfection. Antigen testing in such circumstances is preferred, with the specimen collected within the first five to seven days from symptom onset ([KDHE, 2021](#); [CDC, 2021](#)).

### **Post-exposure management of vaccinated individuals (updated January 2022)**

Exempt from quarantine IF they meet ALL the following criteria:

1. Asymptomatic following their exposure.
2. Up to date for their COVID-19 vaccinations (received all recommended COVID-19 vaccines, including any booster dose(s) when eligible).
  - a. For individuals not eligible for booster doses, they must be at least two weeks out from their last primary dose (second dose of Moderna or Pfizer or single dose of Johnson and Johnson).
  - b. Individuals are eligible to receive a booster dose 5 months after the second dose of Moderna or Pfizer vaccine or 2 months after receiving a single dose of Johnson and Johnson.
3. Exposed vaccinated staff/children who remain in the childcare setting are **strongly recommended** to wear a mask (covering mouth and nose) **at all times**.

Those meeting the above criteria can continue to work/attend childcare unless they become symptomatic. It is recommended by the CDC that fully vaccinated individuals get tested via PCR or antigen test three to five days after exposure to a suspected or confirmed case of COVID-19 even if they do not have symptoms. However, they do not have to isolate at home while waiting for results if they do not have symptoms. Fully vaccinated individuals that do not have symptoms do not need to quarantine but should mask while in public indoor settings for 10 days after exposure.

If the exposed person becomes symptomatic, they should be tested via PCR or antigen test. Receiving the vaccine does not affect the results of a PCR or antigen test.

## **COVID-19 Vaccination**

Getting vaccinated as soon as the opportunity is available is an important way for providers, staff and provides family to stay safe and reduce the risk of getting seriously ill from COVID-19. Review CDC's [COVID-19 Vaccination Information](#) or talk to a healthcare provider for more information. Even after childcare providers and staff are vaccinated, there will be a need to continue prevention measures for the foreseeable future including wearing masks, physical distancing and other important prevention strategies outlined in this guidance document.

### **Additional doses of vaccine for those with moderately to severely compromised immune systems**

Currently, CDC is recommending that moderately to severely immunocompromised people receive an additional dose. This includes people who have:

- Been receiving active cancer treatment for tumors or cancers of the blood.
- Received an organ transplant and are taking medicine to suppress the immune system.
- Received a stem cell transplant within the last 2 years or are taking medicine to suppress the immune system.

- Moderate or severe primary immunodeficiency (such as DiGeorge syndrome, Wiskott-Aldrich syndrome).
- Advanced or untreated HIV infection.
- Active treatment with high-dose corticosteroids or other drugs that may suppress your immune response.

People should talk to their healthcare provider about their medical condition and whether getting an additional dose is appropriate for them. Immunocompromised people should continue to follow prevention measures such as wearing a mask and physical distancing until advised otherwise by their healthcare provider. Close contacts of immunocompromised people should also be encouraged to be vaccinated against COVID-19 to help protect others.

**COVID-19 booster doses**

Individuals age 12 and up are currently recommended to have a booster dose. To see the most current recommendations for booster doses, please visit: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/stay-up-to-date.html>.

**Testing**

**Types of Tests:**

<p><b><u>Antigen OR Rapid Diagnostic Test (RDT):</u></b></p>	<p>Antigen tests detect a protein on the virus. Results for most antigen tests are available onsite in 15-30 minutes. They may be useful as an initial data point, but because antigen tests may not detect lower levels of the virus, false negatives are a concern. If COVID-19 is suspected or there has been a known exposure, an RDT/antigen test should be followed by a confirmatory PCR to make a final diagnosis.</p> <p>At home antigen tests are more reliable when an individual has symptoms and the test is done 24 hours after symptom onset. The manufactures directions must be followed exactly, which could include testing more than once with the home kit. If a person tests positive with any of these methods, they should isolate at home.</p>
<p><b><u>PCR/molecular test:</u></b></p>	<p>Polymerase chain reaction tests detect the presence of viral genetic material in specimens. These tests take longer (sometimes several days) because they must be sent to a lab for processing but are generally more sensitive than antigen tests. JCDHE currently offers free PCR tests (nasal swab version). There are now some rapid PCR tests available as well in the community.</p>

<b><u>Serology:</u></b>	Blood test that detects antibodies one may have to the virus from an immune system response. These are NOT diagnostic tests and should not be used as such. Serology tests do not provide sufficient evidence of immunity and cannot be used to release individuals from quarantine.
-------------------------	--

**Acceptable tests for return to childcare setting**

PCR tests are best for confirmation of COVID-19 infections. Serology tests are not diagnostic tests and, therefore, are never sufficient to prove current infection. Antigen tests (RDTs) are a gray area. Childcare personnel can use **positive** antigen tests as confirmation of a COVID-19 positive individual. Antigen tests present concerns about false negatives; therefore, **negative** antigen tests on symptomatic individuals (1 primary and/or ≥2 secondary symptoms) should NOT be used to return to childcare setting. However, if the person had a previous positive diagnosis in the last 90 days, then test using an antigen test to avoid picking up remnants for previous infection.

Individuals with one primary or two or more secondary symptoms, but negative COVID-19 test result, need a physician’s alternate diagnosis or wait 10 days from symptom onset before returning to childcare. Individuals with a positive antigen test without a subsequent negative PCR test within 48 hours of the initial antigen test will be considered presumed positive. If a negative PCR test is obtained within 48 hours of the initial antigen test, then the individual would not be considered a case.

When PCR tests are not available it is acceptable to do the antigen 2 days in a row – if both are negative the person is not considered positive.

**Exclusion while waiting for results**

Current or recent symptomatic individuals awaiting COVID-19 test results should be excluded from childcare until confirmatory lab results are received. Individuals who are waiting on test results prior to planned travel or a medical procedure do not need to be excluded. Vaccinated staff without symptoms after exposure do not have to quarantine while waiting on results.

**Period of immunity**

Individuals with lab documentation of previous infection no more than 90 days prior to the most recent exposure (or within the CDC’s most recent guidelines) MAY be released from quarantine recommendations.

**Exposures outside of the childcare setting**

If a child or staff member is not up to date for COVID-19 vaccinations or has not had a lab-documented COVID-19 infection in the last 90 days and is a close contact of a positive individual, no matter the setting in which they were exposed, the **safest option** is to exclude them from childcare settings.

All close contacts should self-monitor for symptoms for 14 days from exposure. If symptoms develop during the 14-day period, person should self-isolate/be excluded from childcare setting and get a PCR test.

### **Household contact**

If a household member (sibling, parent, etc.) tests positive for COVID-19, then all other household members must be quarantined per current public health recommendations following their last interaction with the positive case. If the positive individual can isolate in a separate bedroom, with a separate bathroom, spending little to no time in common areas and always wearing a mask in the presence of other household members, then the quarantine begins on the day the positive individual began isolating away from the household. If this is not possible, then household members will need to quarantine per current public health recommendations following the end of the infected person's isolation. This may mean that family members are quarantined for longer periods. If additional household members become symptomatic/test positive during the isolation or quarantine period, the quarantine period starts over.

### **Contacts of contacts**

If an individual is notified that they are a close contact of a COVID-19 positive individual, only that person who was directly exposed needs to quarantine. Other family members (e.g., siblings) do not need to quarantine if they did not have contact with the infected individual.